

## **EFFECT OF DIETARY CALCIUM ON GROWTH PERFORMANCE OF GROWING PIGS**

S. A. Lee, L. V. Lagos, and H. H. Stein  
Division of Nutritional Sciences, University of Illinois  
Urbana, IL 61801, USA

### **1.- INTRODUCTION**

Relative bioavailability of P has been used to evaluate the available P in inorganic P supplements and values have usually been determined relative to the availability of P in monosodium phosphate or monocalcium phosphate [1]. However, only limited data for the bioavailability of Ca in inorganic supplements have been reported [2], but it is believed that the availability of Ca in most Ca supplements is high [3,4]. Values for relative bioavailability of P are variable among sources of P [5], and the relative bioavailability of P is often greater than digestibility values [6]. Therefore, use of values for the digestibility of Ca and P has been suggested as a more accurate way of assessing the digestibility and requirement of Ca and P by pigs [2,7].

### **2.- DIGESTIBILITY OF CA AND P BY GROWING PIGS**

The digestibility of a nutrient represents the amount of that nutrient that disappears from the intestinal tract and it is assumed that this amount is also available for metabolism after absorption [8]. Total tract digestibility is used to determine the digestibility of Ca and

P because there is no net absorption or secretion of Ca and P into the large intestine [7,9]. Apparent total tract digestibility (ATTD) values (%) can be calculated as previously outlined [10] using eq. [1]:

$$\text{ATTD} = \frac{\text{intake} - \text{output}}{\text{intake}} \times 100, \quad [1]$$

where intake and output of nutrients in feces are expressed as g per d.

The values for ATTD are usually influenced by dietary nutrient levels because not only dietary nutrients that have not been digested and absorbed, but also nutrients of endogenous origin are excreted in the fecal output, which may result in an underestimation of ATTD values [11,12]. The endogenous loss of nutrients from pigs consist of the basal endogenous loss that is considered an inevitable loss from the body that is related to dry matter intake (DMI) and a diet specific endogenous loss that is influenced by dietary components [13]. The ATTD values can be corrected for either basal endogenous loss or total endogenous loss to calculate standardized total tract digestibility (STTD) or true total tract digestibility (TTTD), respectively. Because the STTD or TTTD values are not affected by the level of nutrients in the diet, the STTD and TTTD of Ca and P are believed to be additive in mixed diets [14,15].

Basal endogenous losses of Ca and P have been measured by feeding a Ca-free or P-free diet to pigs [16,17] and are calculated as previously outlined (adapted from [10]) using eq. [2]:

$$\text{Basal endogenous loss} = \frac{\text{output of Ca or P}}{\text{DMI}} \times 1,000, \quad [2]$$

where basal endogenous loss is expressed in mg/kg of DM, DMI in kg of DM/d and the fecal output in g/d.

The STTD values (%) can be calculated from the following eq. [3] (adapted from [10]):

$$\text{STTD} = \frac{\text{intake} - (\text{output} - \text{daily basal endogenous loss})}{\text{intake}} \times 100, \quad [3]$$

where intake, output, and daily basal endogenous loss are in g/d.

Values for the basal endogenous loss of P that are estimated using a P-free diets are relatively constant regardless of BW and an average of a number of experiments indicated that a value of 190 mg/kg DMI is representative of the basal endogenous loss of P [2,18]. However, the basal endogenous loss of Ca that is estimated from pigs fed corn-based or cornstarch-based diets appears to be more variable and values ranging from 123 to 550 mg/kg DMI have been reported (Table 1). Relatively lower values for the basal endogenous loss of Ca and the ATTD of Ca have been observed in pigs fed cornstarch-based diets compared with pigs fed corn-based diets [16].

**Table 1.- Estimates of basal endogenous loss and total endogenous loss of Ca by growing pigs**

Reference	Initial BW, kg	Endogenous loss of Ca, mg/kg DMI	Main ingredients
Ca-free diet (basal endogenous loss)			
González-Vega et al. [16]	19.2	220	Cornstarch
González-Vega et al. [16]	19.4	396	Corn
González-Vega et al. [29]	17.7	123	Cornstarch
Merriman, Stein [34]	15.4	329	Corn
Merriman [47]	14.9	550	Corn
Blavi et al. [31]	15.4	430	Corn
Santana et al. [48]	20.5	140	Corn, soybean meal (1.25% soybean meal)
Lee et al. [27]	19.8	430	Corn
Regression method (total endogenous loss)			
Zhang, Adeola [15]	20.0	207	Corn, corn gluten meal, limestone
Zhang, Adeola [15]	20.0	316	Corn, corn gluten meal, dicalcium phosphate
Zhang, Adeola [15]	20.0	264	Corn, corn gluten meal, limestone, dicalcium phosphate
González-Vega et al. [33]	16.7	160	Canola meal
González-Vega et al. [33]	16.7	189 (with phytase)	Canola meal

**2.1.- Absorption of Ca**

The mechanism of Ca absorption is modified based on the quantity of Ca in the diet [19]. Transcellular transport is the primary route under low dietary Ca conditions whereas paracellular transport through the tight junctions is the preferred route of absorption if dietary Ca is at adequate or high levels [19]. This is because of the hormonal regulation of Ca in plasma by the active form of vitamin D (1,25-dihydroxycholecalciferol; calcitriol) and calcitonin [20]. Vitamin D is activated in the kidney by the parathyroid hormone (PTH) which is released as a result of low concentration of Ca in plasma, which increases the transcellular intestinal absorption of Ca and the reabsorption of Ca from the renal tubules and bones [20]. Intestinal and renal Ca absorption is increased by increasing the expression of apical membrane channels (transient receptor potential vanilloid; TRPV), which allow Ca to enter the cell, Ca binding proteins (calbindin), which transport Ca through the cell, and basolateral membrane channels (Ca-ATPase), which allow Ca to exit

the cell [21]. Calcitonin, by contrast, is released in response to high concentrations of plasma Ca and increases the excretion of Ca in urine and preventing the resorption of Ca from bones [20].

## 2.2.- Sources of Ca and P

Most dietary Ca is supplied by mineral supplements, but animal and plant ingredients also provide Ca. Mineral supplements mainly include calcium carbonate and calcium phosphates, and the concentrations of Ca range from 15 to 40% [2]. Cereal grains and co-products of cereal grains, and oilseed meals can also provide dietary Ca, although most of the plant ingredients are low in Ca (range = 0.02 to 1.17%; [2]). Animal feed ingredients including milk products and animal byproducts contain relatively greater concentrations of Ca (between 0.20 and 8.28%) than plant feed ingredients [2]. The ATTD and STTD of Ca in feed ingredients have been evaluated in recent years and summarized values are presented in Table 2.

**Table 2.- Apparent total tract digestibility (ATTD), standardized total tract digestibility (STTD), and true total tract digestibility (TTTD) of Ca in feed ingredients without and with phytase added to the diet fed to growing pigs**

Item, % Supplementation of phytase <sup>1</sup>	ATTD of Ca		STTD of Ca		TTTD of Ca	
	-	+	-	+	-	+
Mineral supplements						
Monocalcium phosphate <sup>2</sup>	83	83	86	86	-	-
Dicalcium phosphate <sup>2, 3</sup>	73	76	77	79	76	-
Calcium carbonate <sup>2, 3, 4, 5, 6, 7, 8, 9</sup>	68	74	71	77	70	-
Calcium carbonate without fat source <sup>6</sup>	52	-	-	-	-	-
Lithothamnium calcareum <sup>2</sup>	63	66	65	69	-	-
Plant feed ingredients						
Canola meal <sup>7, 8, 10</sup>	41	-	45	70	47	70
Soybean meal <sup>7, 8, 11</sup>	53	-	78	-	-	-
Sugar beet co-product <sup>2</sup>	66	63	68	65	-	-
Sunflower meal <sup>8</sup>	22	-	-	-	-	-
Animal feed ingredients						
Meat and bone meal <sup>12</sup>	75	-	77	82	-	-
Meat meal <sup>12</sup>	75	-	77	86	-	-
Fish meal <sup>13</sup>	62	71	65	73	-	-
Poultry meal <sup>12</sup>	85	74	82	76	-	-
Poultry by product meal <sup>12</sup>	81	84	88	87	-	-
Skim milk powder <sup>7</sup>	95	-	97	-	-	-
Whey powder <sup>7</sup>	97	-	99	-	-	-
Whey permeate <sup>7</sup>	61	-	63	-	-	-

<sup>1</sup>Phytase level varies from 500 to 1,500 phytase units/kg diet; <sup>2</sup>González-Vega et al. [29]; <sup>3</sup>Zhang, Adeola [15]; <sup>4</sup>Blavi et al. [31]; <sup>5</sup>Merriman, Stein [34]; <sup>6</sup>Merriman et al. [37]; <sup>7</sup>Unpublished data from the University of Illinois; <sup>8</sup>Zhang et al. [49]; <sup>9</sup>Kwon, Kim [50]; <sup>10</sup>González-Vega et al. [33]; <sup>11</sup>Bohlke et al. [9]; <sup>12</sup>Merriman et al. [51]; <sup>13</sup>González-Vega et al. [16].

### 2.3.- Comparative digestibility values for Ca and P by gestating sows and growing pigs.

The digestibility of energy and some nutrients is affected by the physiological state of the animal and sows usually have greater digestibility values than growing pigs [22,23]. Absorption and retention of Ca and P also increases during pregnancy in humans and rats compared with non-pregnant periods because of an increased need for maternal body and fetus Ca and P [24]. However, pig data have demonstrated that ATTD of Ca and P in growing pigs are greater than in sows [25,26]. Difference in feed intake of gestating sows and growing pigs may affect the ATTD values, but it is not likely that the difference in feed intake is the main reason for the difference because feed intake of sows does not affect digestibility of Ca and P [26]. Greater endogenous losses from gestating sows than from growing pigs may also affect the ATTD values. Basal endogenous loss of Ca from gestating sows fed a Ca-free diet was 1,580 mg/kg DMI, whereas it was 430 mg/kg DMI for growing pigs (Table 3; [27]). However, gestating sows had reduced values for STTD of Ca and P compared with growing pigs indicating that the differences in endogenous losses were not the main reason for the differences in digestibility between growing pigs and sows. Considering the fact that retention of Ca and P is low in gestating sows, it is possible that gestating sows have a requirement for Ca and P that is close to the maintenance requirement, whereas growing pigs have a requirement for growth and bone development in addition to the requirement for maintenance, but additional research is needed to address this hypothesis.

**Table 3.- Apparent total tract digestibility (ATTD) of DM and standardized total tract digestibility (STTD) of Ca and P by sows in mid-gestation and growing pigs fed diets containing different levels of phytate<sup>1</sup>**

Item	Physiological state	Gestating sows		Growing pigs	
		0.98	2.94	0.98	2.94
Dietary phytate (%)					
ATTD of DM		90.1	81.6	90.2	80.9
STTD of Ca <sup>2</sup>		25.6	26.2	78.3	44.8
STTD of P <sup>3</sup>		27.8	11.8	63.0	27.3

<sup>1</sup>Derived from [27].

<sup>2</sup>The values for STTD of Ca were calculated by correcting ATTD of Ca for basal endogenous loss of Ca; basal endogenous loss of Ca from gestating sows = 1,580 mg/kg DM intake; basal endogenous loss of Ca from growing pigs = 430 mg/kg DM intake.

<sup>3</sup>The values for STTD of P were calculated by correcting ATTD of P for basal endogenous loss of P; basal endogenous loss of P from gestating sows = 780 mg/kg DM intake; basal endogenous loss of P from growing pigs = 160 mg/kg DM intake.

### 2.4.- Dietary factors affecting digestibility of Ca in pigs

Factors such as phytate [28] and fiber [16] as well as inclusion of phytase [29,30] or Zn [31] in the diet affect the digestibility of Ca. By contrast, factors such as dietary Ca level [7,32,33], particle size of Ca supplements [4,34], and concentration of P [35,36] do not appear to affect the digestibility of Ca. The effect of different sources of fat in diets on the digestibility of Ca is not clear although most fat sources increase Ca digestibility [37].

### 3.- REQUIREMENTS FOR CALCIUM BY GROWING PIGS

The concentration of dietary Ca and P as well as the Ca:P ratio [21], the concentration of vitamin D in the diet [20], and the age of the animal [2] may affect the requirement for Ca by growing pigs. Inclusion of phytase and other additives, animal genetics, concentration of energy in the diet, and management strategies can also influence the requirement for Ca [38].

#### 3.1.- Requirements for Total Ca

Empirical measurements and factorial calculations have been used to determine requirements for total Ca by pigs [20] and although factorial calculations are believed to be more accurate for establishing nutrient requirements, empirical measurements are more often used [39]. Growth performance and bone development are the most common response criteria for the empirical method, however, blood composition and carcass characteristics have also been used [40,41]. In these type of experiments a constant concentration of P and different concentrations of Ca or a constant Ca:P ratio may be used. Results have demonstrated that the quantities of Ca and P needed to maximize bone development are greater than what is needed to maximize growth performance.

Because of the interest in reducing P excretion in the manure, maximizing P retention, and maximizing growth performance of pigs, values for STTD P are used in diet formulation [2]. However, requirements for Ca have been expressed on the basis of total Ca due to a lack of data about the digestibility of Ca in Ca containing feed ingredients [2]. A factorial approach was used to calculate the requirement for both STTD P and total Ca, thus, the STTD P requirement was based on maximizing whole-body P retention and the requirement for total Ca was calculated by multiplying the STTD P requirement by 2.15 [2]. However, the basis for this value is unclear. The STTD P requirement (%) was obtained using eq. [5]:

$$\text{STTD P requirement} = 0.85 \times \left[ \frac{(\text{maximum whole-body P retention})}{0.77} + 0.19 \times \text{DMI} + 0.007 \times \text{BW} \right], [5]$$

where it was assumed that 1) 85% of P requirement for maximum P retention will maximize growth performance; 2) P body mass is directly related to body protein mass; 3) 77% of the STTD P intake is used for P retention; 4) 190 mg/kg DMI are estimated as endogenous losses of P; and 5) 7 mg/kg BW are considered the minimum P urine losses. The body P mass (g) was obtained by using eq. [6]:

$$\text{Body P mass} = 1.1613 + 26.012 \times \text{body protein mass} + 0.2299 \times (\text{body protein mass})^2, [6]$$

### 3.2.- Requirements for Digestible Ca

Requirements for Ca by pigs will be more accurately estimated using a ratio between STTD Ca and STTD P. Therefore, studies have been conducted to generate values for the STTD of Ca in most Ca containing feed ingredients [42], which allowed for determining the requirements for STTD Ca by growing pigs. Thus, a number of studies have been conducted to establish the requirement for STTD Ca expressed as a ratio between STTD Ca and STTD P.

In a 22-d study, a fixed concentration of STTD P, and 6 concentrations of STTD Ca were used to determine the amount of STTD Ca required to maximize growth performance, bone ash, and Ca retention in 11- to 25-kg pigs [43]. Results indicated that bone ash and Ca retention are maximized at STTD Ca concentrations of minimum 0.48% and 0.60%, respectively. However, a negative effect of STTD Ca above 0.50% on ADG and G:F prevented the estimation of a STTD Ca concentration that maximized growth performance. Therefore, the following studies were conducted using different concentrations of STTD P that were below, at, and above the requirement to estimate STTD Ca requirements an optimal STTD Ca:STTD P ratios at different concentrations of STTD P.

Four experiments have been conducted using 5 levels of STTD Ca and 3 or 4 levels of STTD P [44-46], for a total of 15 or 20 different STTD Ca:STTD P ratios. In the first experiment, the requirement for STTD Ca by 25- to 50-kg pigs was estimated by using 5 and 4 dietary concentrations of STTD Ca and STTD P, respectively. Results indicated a negative effect of increasing dietary concentrations of STTD Ca on growth performance and demonstrated that the impact is more evident if STTD P is at or below the requirement than if P is above the requirement [44], which demonstrates the importance of formulating diets based on a ratio between STTD Ca and STTD P. Growth performance and bone ash were maximized at STTD Ca:STTD P ratios of 1.16:1 to 1.43:1 and 1.55:1 to 1.77:1, respectively.

Two experiments were conducted to determine the requirement for STTD Ca by 100- to 130-kg and 50- to 85-kg pigs [45,46] using a total of 15 different STTD Ca:STTD P ratios. These experiments confirmed that excess concentrations of STTD Ca are detrimental to growth performance but that the negative effect may be ameliorated if dietary STTD P is included above the requirement. Linear responses in the 100- to 130-kg pigs study prevented estimation of maximum responses, however, data indicated greater values for growth performance and bone ash at a ratio between STTD Ca and STTD P below 1.10:1 and above 2.30:1, respectively, if STTD P was at the requirement [45]. For the 50- to 85-kg pigs study, bone ash was maximized at a ratio between STTD Ca and STTD P that ranged from 1.59:1 to 2.03:1, whereas growth performance was maximized at a ratio below 1.25:1, if STTD P was at the requirement [46].

Five levels of STTD Ca and 4 levels of STTD P were used to determine the STTD Ca requirement of 11- to 25-kg pigs (unpublished data). Results were in agreement with



the previous experiments and indicated that maximizing growth performance requires an STTD Ca:STTD P ratio below 1.40:1, if STTD P was at the requirement. At the same STTD P level, bone ash was maximized at a ratio between STTD Ca and STTD P of 1.66:1.

#### 4.- CONCLUSIONS

Growth performance of pigs is negatively affected by increasing dietary concentrations of STTD Ca if dietary P is at or below the requirement. However, the effect is ameliorated if P is above the requirement, indicating that diets should be formulated based on a ratio between STTD Ca and STTD P. Results confirmed that Ca and P requirements for bone ash are greater than for growth performance, which indicates that after the requirement for growth performance is met, pigs are able to utilize Ca and P for bone synthesis. It was also demonstrated that the requirement for Ca expressed as STTD Ca:STTD P that maximizes bone development increases as pig body weight increases, whereas the STTD Ca:STTD P ratio that maximizes growth performance is reduced as the weight of the pig increases (Table 4).

**Table 4.- Requirements for Ca to maximize growth performance and bone ash expressed as a ratio between standardized total tract digestible (STTD) Ca and STTD P for growing and finishing pigs**

Item	Body weight range, kg			
	11 to 25	25 to 50	50 to 85	100 to 130
Growth performance	< 1.40:1	< 1.35:1	< 1.25:1	< 1.10:1
Bone ash	1.70:1	1.80:1	2.00:1	2.30:1

#### 5.- ABSTRACT

Values for standardized total tract digestibility (STTD) are believed to be additive in a complete diet. Therefore, use of values for STTD of Ca and P results in the most accurate diet formulations and values for the STTD of Ca in Ca supplements and feed ingredients have been reported. By supplementing exogenous phytase to swine diets, the STTD of Ca in feed ingredients may increase and, therefore, the STTD values without or with phytase are also available. Values for the STTD of Ca and P by sows are reduced compared with growing pigs, indicating that formulating diets for sows using the STTD values that were obtained using growing pigs may result in inaccuracies. Based on the STTD Ca values obtained from growing pigs, experiments have been conducted to determine the requirement for STTD Ca by growing pigs in recent years. Results of the experiments have demonstrated that increasing concentrations of dietary Ca reduces feed intake and, therefore, reduces average daily gain of pigs. The negative effects of excess Ca



on growth performance is more pronounced in diets with marginal P, but it may be ameliorated by increasing dietary P above the requirement. Therefore, it was concluded that the ratio between STTD Ca and STTD P is very important to optimize growth performance of pigs. The estimated ratios between STTD Ca and STTD P at different weight groups of pigs indicated that the STTD Ca:STTD P ratio to maximize growth performance decreases from 1.40:1 to 1.10:1, as pig body weight increases from 11 to 130 kg, whereas the STTD Ca:STTD P ratio to maximize bone ash increases from 1.70:1 to 2.30:1, as the weight of the pig increases. Results also demonstrated that to optimize growth performance of pigs, less quantities of dietary Ca are needed compared to what is needed to maximize bone ash. It is, therefore, likely that pigs that are destined for the breeding herd have greater requirements for dietary Ca and dietary P than terminal pigs.

## 6.- REFERENCES

1. NRC. Nutrient requirements of swine. 10th rev. ed. Washington, DC, USA: Natl. Acad. Press; 1998.
2. NRC. Nutrient requirements of swine. 11th rev. ed. Washington, DC, USA: Natl. Acad. Press; 2012.
3. Kuznetsov SG, Kal'nitskii BD, Bataeva AP. Biological availability of calcium from chemical compounds for young pigs. *Soviet Agric Sci.* 1987;3:32-4.
4. Ross RD, Cromwell GL, Stahly TS. Effects of source and particle size on the biological availability of calcium in calcium supplements for growing pigs. *J Anim Sci.* 1984;59:125-34.
5. Petersen GI, Pedersen C, Lindemann MD, Stein HH. Relative bioavailability of phosphorus in inorganic phosphorus sources fed to growing pigs. *J Anim Sci.* 2011;89:460-6.
6. Baker SR, Kim BG, Stein HH. Comparison of values for standardized total tract digestibility and relative bioavailability of phosphorus in dicalcium phosphate and distillers dried grains with solubles fed to growing pigs. *J Anim Sci.* 2013;91:203-10.
7. González-Vega JC, Walk CL, Liu Y, Stein HH. The site of net absorption of Ca from the intestinal tract of growing pigs and effect of phytic acid, Ca level and Ca source on Ca digestibility. *Arch Anim Nutr.* 2014;68:126-42.
8. Stein HH. Procedures for determining digestibility of amino acids, lipids, starch, fibre, phosphorus, and calcium in feed ingredients fed to pigs. *Anim Prod Sci.* 2017;57:2317-24.
9. Bohlke RA, Thaler RC, Stein HH. Calcium, phosphorus, and amino acid digestibility in low-phytate corn, normal corn, and soybean meal by growing pigs. *J Anim Sci.* 2005;83:2396-403.
10. Almeida FN, Stein HH. Performance and phosphorus balance of pigs fed diets formulated on the basis of values for standardized total tract digestibility of phosphorus. *J Anim Sci.* 2010;88:2968-77.

11. Fan MZ, Archbold T, Sauer WC, Lackeyram D, Rideout T, Gao YX et al. Novel methodology allows simultaneous measurement of true phosphorus digestibility and the gastrointestinal endogenous phosphorus outputs in studies with pigs. *J Nutr.* 2001;131:2388-96.
12. Zhai H, Adeola O. True total-tract digestibility of phosphorus in monocalcium phosphate for 15-kg pigs. *J Anim Sci.* 2012;90:98-100.
13. Stein HH, Sève B, Fuller MF, Moughan PJ, de Lange CFM. Invited review: Amino acid bioavailability and digestibility in pig feed ingredients: Terminology and application. *J Anim Sci.* 2007;85:172-80.
14. She Y, Wang QY, Stein HH, Liu L, Li DF, Zhang S. Additivity of values for phosphorus digestibility in corn, soybean meal, and canola meal in diets fed to growing pigs. *Asian-Australas J Anim Sci.* 2018;31:1301-7.
15. Zhang F, Adeola O. True is more additive than apparent total tract digestibility of calcium in limestone and dicalcium phosphate for twenty-kilogram pigs fed semipurified diets. *J Anim Sci.* 2017;95:5466-73.
16. González-Vega JC, Walk CL, Stein HH. Effect of phytate, microbial phytase, fiber, and soybean oil on calculated values for apparent and standardized total tract digestibility of calcium and apparent total tract digestibility of phosphorus in fish meal fed to growing pigs. *J Anim Sci.* 2015;93:4808-18.
17. Petersen GI, Stein HH. Novel procedure for estimating endogenous losses and measurement of apparent and true digestibility of phosphorus by growing pigs. *J Anim Sci.* 2006;84:2126-32.
18. Son AR. Effects of dietary fiber on digestibility of nutrients and endogenous losses of phosphorus in pigs. PhD. Diss. Konkuk University, Seoul, Republic of Korea; 2016.
19. Bronner F. Mechanisms of intestinal calcium absorption. *J Cell Biochem.* 2003;88:387-93.
20. Crenshaw TD. Calcium, phosphorus, vitamin D, and vitamin K in swine nutrition. In: *Swine Nutrition*. Edited by Lewis AJ, Southern LL. 2nd ed. Boca Raton, FL: CRC Press 2001. p. 187-212.
21. Veum TL. Phosphorus and calcium nutrition and metabolism. In: *Phosphorus and calcium utilization and requirements in farm animals*. Edited by Vitti DMSS, Kebreab E. Wallingford, UK: CABI 2010. p. 94-111.
22. Casas GA, Stein HH. Gestating sows have greater digestibility of energy in full fat rice bran and defatted rice bran than growing gilts regardless of level of feed intake. *J Anim Sci.* 2017;95:3136-42.
23. Le Goff G, Noblet J. Comparative total tract digestibility of dietary energy and nutrients in growing pigs and adult sows. *J Anim Sci.* 2001;79:2418-27.
24. Kovacs CS. Maternal mineral and bone metabolism during pregnancy, lactation, and post-weaning recovery. *Physiol Rev.* 2016;96:449-547.
25. Kemme PA, Jongbloed AW, Mroz Z, Beynen AC. The efficacy of *Aspergillus niger* phytase in rendering phytate phosphorus available for absorption in pigs is influenced by pig physiological status. *J Anim Sci.* 1997;75:2129-38.

26. Lee SA, Casas GA, Stein HH. The level of feed intake does not influence digestibility of calcium and phosphorus in diets fed to gestating sows, but gestating sows have reduced digestibility of calcium and phosphorus compared with growing gilts. *Can J Anim Sci.* 2018;98:591-4.
27. Lee SA, Walk CL, Stein HH. Comparative digestibility and retention of calcium and phosphorus by gestating sows and growing pigs fed low- and high-phytate diets without or with microbial phytase. *J Anim Sci.* 2018;96(Suppl. 2):83 (Abstr.).
28. Almaguer BL, Sulabo RC, Liu Y, Stein HH. Standardized total tract digestibility of phosphorus in copra meal, palm kernel expellers, palm kernel meal, and soybean meal fed to growing pigs. *J Anim Sci.* 2014;92:2473-80.
29. González-Vega JC, Walk CL, Stein HH. Effects of microbial phytase on apparent and standardized total tract digestibility of calcium in calcium supplements fed to growing pigs. *J Anim Sci.* 2015;93:2255-64.
30. Rodríguez DA, Sulabo RC, González-Vega JC, Stein HH. Energy concentration and phosphorus digestibility in canola, cottonseed, and sunflower products fed to growing pigs. *Can J Anim Sci.* 2013;93:493-503.
31. Blavi L, Sola-Oriol D, Perez JF, Stein HH. Effects of zinc oxide and microbial phytase on digestibility of calcium and phosphorus in maize-based diets fed to growing pigs. *J Anim Sci.* 2017;95:847-54.
32. Stein HH, Adeola O, Cromwell GL, Kim SW, Mahan DC, Miller PS. Concentration of dietary calcium supplied by calcium carbonate does not affect the apparent total tract digestibility of calcium, but decreases digestibility of phosphorus by growing pigs. *J Anim Sci.* 2011;89:2139-44.
33. González-Vega JC, Walk CL, Liu Y, Stein HH. Endogenous intestinal losses of calcium and true total tract digestibility of calcium in canola meal fed to growing pigs. *J Anim Sci.* 2013;91:4807-16.
34. Merriman LA, Stein HH. Particle size of calcium carbonate does not affect apparent and standardized total tract digestibility of calcium, retention of calcium, or growth performance of growing pigs. *J Anim Sci.* 2016;94:3844-50.
35. Stein HH, Boersma MG, Pedersen C. Apparent and true total tract digestibility of phosphorus in field peas (*Pisum sativum* L.) by growing pigs. *Can J Anim Sci.* 2006;86:523-5.
36. Stein HH, Kadzere CT, Kim SW, Miller PS. Influence of dietary phosphorus concentration on the digestibility of phosphorus in monocalcium phosphate by growing pigs. *J Anim Sci.* 2008;86:1861-7.
37. Merriman LA, Walk CL, Parsons CM, Stein HH. Effects of tallow, choice white grease, palm oil, corn oil, or soybean oil on apparent total tract digestibility of minerals in diets fed to growing pigs. *J Anim Sci.* 2016;94:4231-8.
38. Suttle NF. Mineral nutrition of livestock. 4th ed. Wallingford, UK CAB International. 2010.
39. Weremko D, Fandrejowski H, Zebrowska T, Han IK, Kim JH, Cho WT. Bioavailability of phosphorus in feeds of plant origin for pigs - Review. *Asian-Australas J Anim Sci.* 1997;10:551-66.

40. Cromwell GL, Hays VW, Chaney CH, Overfield JR. Effects of dietary phosphorus and calcium level on performance, bone mineralization and carcass characteristics of swine. *J Anim Sci.* 1970;30:519-25.
41. Stockland WL, Blaylock LG. Influence of dietary calcium and phosphorus levels on the performance and bone characteristics of growing-finishing swine. *J Anim Sci.* 1973;37:906-12.
42. Stein HH, Merriman LA, González-Vega JC. Establishing a digestible calcium requirement for pigs. In: *Phytate destruction - consequences for precision animal nutrition.* Edited by Walk CL, Kühn I, Stein HH, Kidd MT, Rodehutschord M. Wageningen, The Netherlands: Wageningen Academic Publishers 2016. p. 207-16.
43. González-Vega JC, Liu Y, McCann JC, Walk CL, Loo JJ, Stein HH. Requirement for digestible calcium by eleven- to twenty-five-kilogram pigs as determined by growth performance, bone ash concentration, calcium and phosphorus balances, and expression of genes involved in transport of calcium in intestinal and kidney cells. *J Anim Sci.* 2016;94:3321-34.
44. González-Vega JC, Walk CL, Murphy MR, Stein HH. Requirement for digestible calcium by 25 to 50 kg pigs at different dietary concentrations of phosphorus as indicated by growth performance, bone ash concentration, and calcium and phosphorus balances. *J Anim Sci.* 2016;94:5272-85.
45. Merriman LA, Walk CL, Murphy MR, Parsons CM, Stein HH. Inclusion of excess dietary calcium in diets for 100- to 130-kg growing pigs reduces feed intake and daily gain if dietary phosphorus is at or below the requirement. *J Anim Sci.* 2017;95:5439-46.
46. Lagos LV, Walk CL, Stein HH. 244 Requirement for digestible calcium at different dietary concentrations of digestible phosphorus indicated by growth performance and bone ash of 50- to 85 kg pigs. *J Anim Sci.* 2018;96(suppl. 2):130-1 (Abstr.).
47. Merriman LA. Factors affecting the digestibility of calcium in feed ingredients and requirements for digestible calcium by pigs. Dissertation. University of Illinois at Urbana-Champaign, Urbana; 2016.
48. Santana ALA, Carvalho PLD, de Oliveira NTE, Gonçalves AC, Gazola AP, Castro DEDE et al. Different sources of calcium for starter pig diets. *Livest Sci.* 2017;206:175-81.
49. Zhang F, Ragland D, Adeola O. Comparison of apparent ileal and total tract digestibility of calcium in calcium sources for pigs. *Can J Anim Sci.* 2016;96:563-9.
50. Kwon WB, Kim BG. Standardized total tract digestibility of phosphorus in various inorganic phosphates fed to growing pigs. *Anim Sci J.* 2017;88:918-24.
51. Merriman LA, Walk CL, Stein HH. The effect of microbial phytase on the apparent and standardized total tract digestibility of calcium in feed ingredients of animal origin. *J Anim Sci.* 2016;94(E-Suppl. 5):110.